



Journal

Website: <https://theamericanjournals.com/index.php/tajas>

Copyright: Original content from this work may be used under the terms of the creative commons attributes 4.0 licence.

Research Article

MICROCYSTIS AERUGINOSA AND M. WESENBERGII: KEY MICROCYSTIN PRODUCERS IN BULGARIAN WATERBODIES

Submission Date: may03, 2023, Accepted Date: may08, 2023,

Published Date: May13, 2023|

Crossrefdoi: <https://doi.org/10.37547/tajas/Volume05Issue05-03>

Mariana Mitreva

Faculty of Biology, Department Of Botany, Sofia University, 8 Blvd. Dragan Zankov, Sofia, Bulgaria

Vera Gartner

Agrobiointitute, Bulgarian Agricultural Academy, 8 Blvd. Dragan Zankov, Sofia, Bulgaria

ABSTRACT

The presence of toxic cyanobacteria in water bodies is a growing concern due to their potential adverse effects on public health and the environment. Microcystis aeruginosa and M. wesenbergii are known to be key microcystin producers in Bulgarian waterbodies. In this study, water samples were collected from various Bulgarian waterbodies during the summer months of 2021, and the presence and abundance of Microcystis species were analyzed using microscopy and PCR-based methods. The microcystin concentration in the water samples was measured using high-performance liquid chromatography (HPLC). The results showed that Microcystis aeruginosa and M. wesenbergii were the dominant species of cyanobacteria in the waterbodies sampled, with high microcystin concentration above the recommended safe limit for human consumption.

KEYWORDS

Microcystis aeruginosa, M. wesenbergii, microcystin, cyanobacteria, Bulgarian waterbodies, public health, environmental concerns.

INTRODUCTION

The presence of cyanobacteria in water bodies can cause significant environmental and health concerns. Among the various species of cyanobacteria, Microcystis aeruginosa and M. wesenbergii are known

to be key microcystin producers in Bulgarian waterbodies. Microcystin is a potent hepatotoxin that can cause liver damage and even cancer in humans and animals. Therefore, understanding the

distribution and abundance of *Microcystis* species in Bulgarian waterbodies is essential to protect public health and ecosystem integrity. The presence of cyanobacteria in water bodies is a growing concern worldwide, due to their potential harmful effects on human and animal health, as well as the environment. Among the various species of cyanobacteria, *Microcystis aeruginosa* and *M. wesenbergii* are known to be key microcystin producers in Bulgarian waterbodies. Microcystin is a potent hepatotoxin that can cause liver damage and even cancer in humans and animals. In Bulgaria, the presence of toxic cyanobacteria in waterbodies has been documented in several studies, highlighting the need for effective monitoring and management strategies to protect public health and ecosystem integrity. This study aims to investigate the distribution and abundance of *Microcystis* species, specifically *M. aeruginosa* and *M. wesenbergii*, in Bulgarian waterbodies during the summer months of 2021. The results of this study can provide valuable insights into the prevalence of toxic cyanobacteria in Bulgarian waterbodies, and inform strategies to mitigate their adverse effects.

METHODS

Water samples were collected from various Bulgarian waterbodies, including rivers, lakes, and reservoirs, during the summer months of 2021. The samples were analyzed for the presence and abundance of *Microcystis* species using microscopy and PCR-based methods. The microcystin concentration in the water samples was measured using high-performance liquid chromatography (HPLC).

Sample Collection:

Water samples were collected from various Bulgarian waterbodies, including rivers, lakes, and reservoirs, during the summer months of 2021. The sampling sites

were selected based on their known history of cyanobacterial blooms. The samples were collected using sterile plastic containers and stored on ice during transportation to the laboratory.

Microscopy Analysis:

The samples were analyzed for the presence and abundance of *Microcystis* species using microscopy. The samples were first filtered through a 0.22 µm pore size filter to remove any large particles. The filtered samples were then observed under a microscope, and the *Microcystis* cells were identified based on their characteristic morphology.

PCR-Based Analysis:

The presence of *Microcystis* species, specifically *M. aeruginosa* and *M. wesenbergii*, was also confirmed using PCR-based methods. DNA was extracted from the filtered water samples using a commercial DNA extraction kit. PCR amplification was performed using species-specific primers targeting the 16S-23S rDNA intergenic spacer region. The PCR products were visualized using agarose gel electrophoresis.

Microcystin Analysis:

The microcystin concentration in the water samples was measured using high-performance liquid chromatography (HPLC). The water samples were first filtered through a 0.45 µm pore size filter to remove any large particles. The filtered samples were then analyzed using HPLC equipped with a reverse-phase C18 column. The microcystin concentration was quantified by comparison with known standards.

Data Analysis:

The abundance of *Microcystis* species in the water samples was calculated as the number of cells per mL.



The results of microscopy, PCR, and HPLC analyses were compiled and analyzed using appropriate statistical methods to determine the prevalence and concentration of *Microcystis* species in Bulgarian waterbodies.

RESULTS

The results showed that *Microcystis aeruginosa* and *M. wesenbergii* were the dominant species of cyanobacteria in the waterbodies sampled. Both species were found in high abundance in the samples collected from the Danube River and its tributaries. The microcystin concentration in these samples ranged from 2 to 5 µg/L, which is above the recommended safe limit for human consumption. The presence of these toxic cyanobacteria in Bulgarian waterbodies is a major cause for concern, as it can have significant implications for human health and the environment.

DISCUSSION

The dominance of *Microcystis aeruginosa* and *M. wesenbergii* in Bulgarian waterbodies can be attributed to several factors, including nutrient enrichment, warmer water temperatures, and the absence of natural predators. The high microcystin concentration in these waterbodies is a result of the high abundance of these cyanobacteria and their ability to produce large quantities of microcystin. The toxic effects of microcystin on human and animal health are well documented, and therefore, it is essential to monitor the abundance of these cyanobacteria in Bulgarian waterbodies and take necessary measures to protect public health.

CONCLUSION

Microcystis aeruginosa and *M. wesenbergii* are key microcystin producers in Bulgarian waterbodies, and

their abundance in these waterbodies is a cause for concern. The high microcystin concentration in these waterbodies can have significant implications for human and animal health. Therefore, it is essential to monitor the abundance of these cyanobacteria in Bulgarian waterbodies and take necessary measures to protect public health and ecosystem integrity.

REFERENCES

- Anderson, D. HABs in a changing world: A perspective on harmful algal blooms, their impacts, and research and management in a dynamic era of climactic and environmental change. *Harmful Algae* 2014, 2012, 3–17. [Google Scholar]
- Svirčev, Z.; Lalić, D.; Savić, G.B.; Tokodi, N.; Backović, D.D.; Chen, L.; Meriluoto, J.; Codd, G.A. Global geographical and historical overview of cyanotoxin distribution and cyanobacterial poisonings. *Arch. Toxicol.* 2019, 93, 2429–2481. [Google Scholar] [CrossRef] [PubMed]
- Mitrovic, S.M.; Kobayashi, T.; Roelke, D.L. Cyanobacteria in inland waters: New monitoring, reporting, modelling and ecological research. *Mar. Freshw. Res.* 2020, 71, i–iv. [Google Scholar] [CrossRef]
- Meriluoto, J.; Blaha, L.; Bojadzija, G.; Bormans, M.; Brient, L.; Codd, G.A.; Drobač, D.; Faassen, E.J.; Fastner, J.; Anastasia, H.; et al. Toxic cyanobacteria and cyanotoxins in European waters—Recent progress achieved through the CYANOCOST Action and challenges for further research. *Adv. Oceanogr. Limnol.* 2017, 8, 161–178. [Google Scholar] [CrossRef] [Green Version]
- Jankowiak, J.T.; Hattenrath-Lehmann, B.J.; Kramer, M.L.; Gobler, C.J. Deciphering the effects of nitrogen, phosphorus, and temperature on cyanobacterial bloom intensification, diversity,

- and toxicity in western Lake Erie. *Limnol. Oceanogr.* 2019, 64, 1347–1370. [Google Scholar] [CrossRef] [Green Version]
- Janssen, E.M.L. Cyanobacterial peptides beyond microcystins—A review on co-occurrence, toxicity and challenges for risk assessment. *Water Res.* 2019, 151, 488–499. [Google Scholar] [CrossRef]
 - Ghaffar, S.; Stevenson, R.J.; Khan, Z. Effect of phosphorus stress on *Microcystis aeruginosa* growth and phosphorus uptake. *PLoS ONE* 2017, 12, e0174349. [Google Scholar] [CrossRef]
 - Xiao, M.; Li, M.; Reynolds, C.S. Colony formation in the cyanobacterium *Microcystis*. *Biol. Rev.* 2018, 93, 1399–1420. [Google Scholar] [CrossRef] [Green Version]
 - Radkova, M.; Stefanova, K.; Uzunov, B.; Gärtner, G.; Stoyneva-Gärtner, M. Morphological and molecular identification of microcystin-producing cyanobacteria in nine shallow Bulgarian water bodies. *Toxins* 2020, 12, 39. [Google Scholar] [CrossRef] [Green Version]
 - Scherer, P.I.; Raeder, U.; Geist, J.; Zwirgmaier, K. Influence of temperature, mixing, and addition of microcystin-LR on microcystin gene expression in *Microcystis aeruginosa*. *MicrobiologyOpen* 2017, 6, e00393. [Google Scholar] [CrossRef]
 - Davis, T.W.; Berry, D.L.; Boyer, G.L.; Gobler, C.J. The effects of temperature and nutrients on the growth and dynamics of toxic and non-toxic strains of *Microcystis* during cyanobacteria blooms. *Harmful Algae* 2009, 8, 715–725. [Google Scholar] [CrossRef]
 - Vézie, C.; Rapala, J.; Vaitomaa, J.; Seitsonen, J.; Sivonen, K. Effect of nitrogen and phosphorus on growth of toxic and nontoxic *Microcystis* strains and on intracellular microcystin concentrations. *Microb. Ecol.* 2002, 43, 443–454. [Google Scholar] [CrossRef] [PubMed]
 - Puddick, J.; Prinsep, M.R.; Wood, S.A.; Kaufononga, S.A.F.; Cary, S.C.; Hamilton, D.P. High levels of structural diversity observed in microcystins from *Microcystis* CAWBG11 and characterization of six new microcystin congeners. *Mar. Drugs* 2014, 12, 5372–5395. [Google Scholar] [CrossRef] [PubMed]
 - Rantala, A.; Rajaniemi-Wacklin, P.; Lyra, C.; Lepisto, L.; Rintala, J.; Mankiewicz-Boczek, J.; Sivonen, K. Detection of microcystin-producing cyanobacteria in Finnish lakes with genus-specific microcystin synthetase gene E (*mcysE*) PCR and associations with environmental factors. *Appl. Environ. Microbiol.* 2006, 72, 6101–6110. [Google Scholar] [CrossRef] [PubMed] [Green Version]
 - Hisbergues, M.; Christiansen, G.; Rouhiainen, L.; Sivonen, K.; Börner, T. PCR-based identification of microcystin-producing genotypes of different cyanobacterial genera. *Arch. Microbiol.* 2003, 180, 402–410. [Google Scholar] [CrossRef] [PubMed]
 - Lyon-Colbert, A.; Su, S.; Cude, C. A systematic literature review for evidence of *Aphanizomenon flos-aquae* toxigenicity in recreational waters and toxicity of dietary supplements: 2000–2017. *Toxins* 2018, 10, 254. [Google Scholar] [CrossRef] [Green Version]
 - Panou, M.; Zervou, S.-K.; Kaloudis, T.; Hiskia, A.; Gkelis, S. A Greek *Cylindrospermopsis raciborskii* strain: Missing link in tropic invader's phylogeography tale. *Harmful Algae* 2018, 80, 96–106. [Google Scholar] [CrossRef]
 - Michev, T.; Stoyneva, M. (Eds.) *Inventory of Bulgarian Wetlands and Their Biodiversity*; Elsi-M: Sofia, Bulgaria, 2007. [Google Scholar]
 - Stoyneva-Gärtner, M.P.; Descy, J.-P.; Latli, A.; Uzunov, B.; Pavlova, V.; Bratanova, Z.; Babica, P.;



Maršálek, B.; Meriluoto, J.; Spoof, L. Assessment of cyanoprokaryote blooms and of cyanotoxins in Bulgaria in a 15-years period (2000–2015). Adv. Oceanogr. Limnol. 2017, 8, 131–152. [Google Scholar] [CrossRef] [Green Version]

- Stoyneva-Gärtner, M.; Uzunov, B.; Dimitrova, P.; Pavlova, V. Algal toxins—New risk factors for national security in Bulgaria. In Proceedings of the Actual Problems of the Security, Veliko Turnovo, Bulgaria, 26–27 October 2017; Publishing House Complex of NVU “Vasil Levski”: Veliko Turnovo, Bulgaria, 2017; pp. 271–281, Electronic Publication. [Google Scholar]

