

## A Clinical Analysis Framework for Assessing the Prevalence and Risk Factors of Incarcerated Inguinal Hernia in Adult Populations

**Dr. Rizky Pratama Santoso**

Department of Oncology and Cancer Research  
Faculty of Medicine, Universitas Airlangga  
Surabaya, Indonesia

**Dr. Nadia Putri Wibowo**

Department of Medical Oncology and Translational Cancer Science  
Faculty of Medicine, Universitas Indonesia  
Jakarta, Indonesia

Received: 25 July 2026 | Received Revised Version: 16 Aug 2026 | Accepted: 10 Sep 2026 | Published: 01 Oct 2026

Volume 08 Issue 10 2026 | DOI:10.37547/tajmspr/Volume08Issue01-07

### Abstract

*Adult inguinal hernia incarceration represents a significant surgical emergency associated with increased morbidity, risk of bowel ischemia, and potential mortality when treatment is delayed. Despite advances in elective hernia repair techniques, incarcerated inguinal hernias continue to present substantial clinical and epidemiological challenges, particularly among elderly patients and individuals with delayed healthcare access. This research and review article adopts an epidemiological approach to evaluate incidence patterns, determinants, and surgical implications of adult inguinal hernia incarceration through a comprehensive synthesis of contemporary literature. The study examines demographic trends, risk factors contributing to incarceration, management algorithms, emergency intervention strategies, and outcomes associated with different surgical approaches. Particular attention is given to factors influencing bowel resection rates and emergency operative decision-making. The findings indicate that age, delayed presentation, hernia duration, comorbidities, and bowel compromise significantly influence clinical outcomes. Contemporary evidence also demonstrates the growing role of manual reduction techniques and laparoscopic surgery in selected emergency settings. The article highlights the importance of early diagnosis, risk stratification, and evidence-based management protocols to reduce complications and optimize patient outcomes. The study contributes to a deeper understanding of incarcerated inguinal hernia epidemiology and provides a framework for improving emergency surgical care.*

**Keywords:** Inguinal Hernia, Hernia Incarceration, Epidemiology, Emergency Surgery, Bowel Resection, Laparoscopic Repair, Taxis Reduction, Surgical Outcomes, Adult Patients, Hernia Management

---

© 2026 Ivan Metelkin. This work is licensed under a Creative Commons Attribution 4.0 International License (CC BY 4.0). The authors retain copyright and allow others to share, adapt, or redistribute the work with proper attribution.

**Cite This Article:** Santoso, D., & Wibowo, D. N. P. (2026). A Clinical Analysis Framework for Assessing the Prevalence and Risk Factors of Incarcerated Inguinal Hernia in Adult Populations. The American Journal of Medical Sciences and Pharmaceutical Research, 8(10), 01–07. Retrieved from <https://theamericanjournals.com/index.php/tajmspr/article/view/8455>

---

## 1. Introduction

Inguinal hernia is among the most frequently encountered surgical conditions worldwide, affecting millions of adults across diverse demographic groups. Although many inguinal hernias remain asymptomatic or minimally symptomatic for extended periods, incarceration represents a severe complication requiring urgent medical attention. Hernia incarceration occurs when herniated tissue becomes trapped within the hernia sac and cannot be reduced back into the abdominal cavity. If left untreated, incarceration may progress to strangulation, bowel ischemia, necrosis, and

systemic complications, necessitating emergency surgical intervention (Pawlak et al., 2021).

The epidemiological significance of incarcerated inguinal hernia lies in its association with increased hospitalization rates, emergency operative procedures, prolonged recovery periods, and higher healthcare expenditures. Adult patients, particularly elderly individuals, are disproportionately affected due to age-related tissue weakness, delayed symptom recognition, and multiple comorbid conditions (Ferranti et al., n.d.). Understanding incidence patterns and determinants is therefore critical for reducing morbidity and improving clinical outcomes.




**Figure 1.** Epidemiological Progression of Adult Inguinal Hernia Incarceration

**Caption:** This figure illustrates the progression pathway from uncomplicated inguinal hernia to incarceration, bowel obstruction, strangulation, and emergency surgical intervention. The model highlights key risk factors including advanced age, delayed treatment, chronic hernia duration, and associated comorbidities. Understanding these progression

stages assists clinicians in identifying high-risk patients and implementing preventive surgical strategies before life-threatening complications occur.

The growing emphasis on evidence-based emergency surgery has led to increased interest in identifying determinants that influence incarceration risk and postoperative outcomes. Several studies have examined predictors of bowel resection, emergency management algorithms, and minimally invasive surgical techniques, creating a foundation for more effective clinical decision-making (Chen et al., 2020).

This article aims to evaluate incidence patterns, epidemiological determinants, and surgical implications associated with adult inguinal hernia incarceration. The study synthesizes available evidence to identify key trends, assess current management strategies, and provide recommendations for improving emergency surgical care.

### Objectives

1. To examine incidence patterns of incarcerated inguinal hernia among adults.
2. To identify determinants associated with incarceration and adverse outcomes.
3. To evaluate emergency management approaches and surgical interventions.
4. To analyze implications for clinical practice and future research.




**Figure 2.** Clinical Management Framework for Incarcerated Inguinal Hernia

**Caption:** The figure presents a clinical decision-making framework for emergency management of incarcerated inguinal hernia. The pathway includes patient assessment, evaluation for strangulation, manual reduction (Taxis), imaging support, laparoscopic or open surgical intervention, and postoperative monitoring. The framework emphasizes timely intervention and risk-based treatment selection to minimize bowel resection and postoperative complications.

### 2. Literature Review

The contemporary literature demonstrates considerable advancement in understanding incarcerated inguinal hernia epidemiology and management. However,

significant variations remain regarding optimal intervention timing, surgical technique selection, and risk prediction models.

Akay and Akici (2020) investigated the role of laparoscopic repair in incarcerated inguinal hernias and argued that minimally invasive approaches may offer substantial benefits, including enhanced visualization, reduced postoperative pain, and shorter hospital stays. Their findings suggest that laparoscopic surgery is increasingly becoming a viable first-line option in selected emergency cases.

Chen et al. (2020) conducted a meta-analysis examining risk factors associated with bowel resection among patients presenting with incarcerated groin hernias. Their study identified delayed presentation, advanced age, and bowel compromise as major predictors of bowel resection. Importantly, the authors emphasized that early intervention significantly reduces adverse outcomes and postoperative complications (Chen et al., 2020).

East et al. (2020) revisited the technique of Taxis, or manual reduction under analgesia and sedation, particularly during the COVID-19 pandemic when reducing emergency surgical burden became essential. Their review demonstrated that appropriately selected patients may safely undergo manual reduction, potentially avoiding immediate surgery and reducing healthcare resource utilization.

Ferranti et al. highlighted unique challenges associated with incarcerated inguinal hernia in elderly populations. Age-related physiological changes, frailty, and coexisting diseases contribute to increased perioperative risks and necessitate individualized management strategies.

Merani et al. (2021) expanded the understanding of unusual incarceration presentations by documenting transplant ureter incarceration within inguinal hernias. Their work demonstrated the diverse anatomical complexities that can influence diagnostic and therapeutic decisions.

Pawlak et al. (2021) proposed an evidence-based algorithm for managing incarcerated inguinal hernia in emergency settings. Their framework integrates patient assessment, Taxis application, imaging evaluation, and surgical decision-making to optimize patient safety and treatment effectiveness.

Sabra et al. (2020) presented a rare case involving perforated sigmoid colon cancer manifesting as an incarcerated inguinal hernia. This study underscored the

importance of differential diagnosis and comprehensive intraoperative evaluation.

Sartori et al. (2023) conducted a systematic review and meta-analysis examining laparoscopic management of acute incarcerated groin hernias. Their findings suggested favorable outcomes associated with minimally invasive approaches, including reduced wound complications and improved postoperative recovery.

### Research Gap

Although existing literature provides substantial insight into emergency management and surgical techniques, limited emphasis has been placed on integrating epidemiological determinants with surgical outcome prediction. Additionally, there remains a need for comprehensive frameworks that combine incidence patterns, risk stratification, and treatment optimization within a unified analytical model.

### 3. Methodology

This study adopts a research-review methodology based exclusively on the provided literature sources. A qualitative epidemiological synthesis approach was employed to examine incidence patterns, determinants, management strategies, and surgical implications of adult inguinal hernia incarceration.

### Conceptual Framework

The analytical framework consists of four interconnected domains:

1. Epidemiological Characteristics
2. Risk Determinants
3. Clinical Management Strategies
4. Surgical Outcomes

### Epidemiological Assessment

The epidemiological component evaluates incidence-related observations reported across studies. Factors considered include age distribution, sex predominance, disease progression, emergency presentation patterns, and complication frequency.

### Determinant Analysis

Determinants influencing incarceration were categorized into:

- Patient-related factors
- Hernia-related factors
- Healthcare access factors
- Surgical timing factors

Patient-related determinants include age, comorbidities, and physiological reserve. Hernia-related determinants encompass duration, size, and reducibility status. Healthcare-related factors involve delayed diagnosis and treatment accessibility.

### Management Evaluation

Emergency management strategies were examined according to:

- Manual reduction techniques
- Open surgical repair
- Laparoscopic repair
- Hybrid approaches

The safety and effectiveness of each strategy were analyzed within the context of available evidence.

### Outcome Assessment

Clinical outcomes evaluated include:

- Bowel resection rates
- Postoperative complications
- Length of hospitalization
- Recovery outcomes
- Mortality risk

Particular attention was given to factors associated with bowel resection because of their strong prognostic significance (Chen et al., 2020).

## 4. Results / Findings

The synthesized evidence reveals several significant epidemiological patterns regarding adult inguinal hernia incarceration.

First, increasing age emerged as one of the most consistent determinants of incarceration and adverse surgical outcomes. Elderly patients exhibited higher rates of emergency presentation, bowel compromise, and postoperative complications. Age-related tissue degeneration and delayed symptom recognition appear to contribute substantially to this trend.

Second, delayed medical presentation significantly increased the likelihood of bowel ischemia and bowel resection. Multiple studies demonstrated that prolonged incarceration duration was strongly associated with tissue compromise and complex surgical intervention requirements. The meta-analysis by Chen et al. (2020) repeatedly identified delayed treatment as a major predictor of bowel resection risk.

Third, manual reduction through Taxis demonstrated effectiveness in carefully selected patients without signs of strangulation. Evidence suggests that this approach can reduce immediate surgical burden while maintaining patient safety when performed under appropriate clinical supervision (East et al., 2020; Pawlak et al., 2021).

Fourth, laparoscopic techniques demonstrated promising outcomes compared with traditional open surgery in selected emergency cases. Improved visualization, reduced wound complications, and shorter recovery periods were consistently reported advantages (Akay&Akici, 2020; Sartori et al., 2023).

The review also identified significant variability in management practices across institutions. Differences in surgeon experience, patient selection criteria, and resource availability influenced treatment decisions and clinical outcomes.

Importantly, bowel resection emerged as a critical outcome indicator. Studies consistently showed that patients requiring bowel resection experienced longer hospitalization, increased postoperative morbidity, and greater healthcare utilization. The risk factors identified by Chen et al. (2020) therefore represent essential considerations in emergency surgical planning.

## 5. Discussion

The findings demonstrate that adult inguinal hernia incarceration remains a clinically significant emergency condition despite advancements in elective hernia management. The epidemiological patterns identified across studies indicate that both biological and healthcare-system factors contribute to disease progression and outcomes.

One of the most important observations is the relationship between delayed intervention and adverse outcomes. The repeated association between treatment delay and bowel resection risk supports the need for proactive surgical evaluation and timely intervention. The findings of Chen et al. (2020) reinforce the concept that early recognition and treatment represent the most effective strategies for reducing severe complications. The consistency of these findings across studies strengthens their clinical relevance.

The increasing adoption of Taxis represents an important evolution in emergency hernia management. Historically, emergency surgery was considered the default approach for incarcerated hernias. However, contemporary evidence indicates that carefully selected patients may benefit from manual reduction, reducing surgical burden while maintaining safety. Nevertheless, inappropriate application may delay necessary surgery and increase risk. Therefore, strict patient selection criteria remain essential.

The growing role of laparoscopic surgery reflects broader trends in minimally invasive surgical practice. Evidence from Akay and Akici (2020) and Sartori et al. (2023) suggests that laparoscopic approaches may improve outcomes by reducing tissue trauma and enhancing intraoperative assessment. However, successful implementation requires specialized expertise and institutional resources.

The literature also highlights the complexity of atypical incarceration presentations. Cases involving transplant ureters or occult malignancies demonstrate that incarceration is not always a straightforward mechanical process. These findings emphasize the importance of comprehensive diagnostic evaluation and individualized treatment planning.

From a theoretical perspective, incarcerated inguinal hernia can be understood as the interaction of anatomical vulnerability, physiological deterioration, and healthcare access factors. This multidimensional framework explains why some patients progress rapidly

to severe complications while others experience more favorable outcomes.

Several limitations were identified within the available literature. Many studies involved retrospective designs, limiting causal interpretation. Variability in patient populations, surgical protocols, and outcome definitions also complicates direct comparison. Furthermore, standardized epidemiological reporting remains limited across regions.

Despite these limitations, the evidence consistently supports risk-based management strategies. The repeated identification of advanced age, delayed presentation, and bowel compromise as adverse prognostic indicators suggests that these variables should be incorporated into future predictive models. Such models could assist clinicians in prioritizing interventions and improving resource allocation.

## 6. Conclusion

Adult inguinal hernia incarceration remains a significant surgical emergency associated with substantial morbidity and healthcare burden. This review demonstrates that incidence patterns and clinical outcomes are strongly influenced by age, treatment delays, hernia characteristics, and comorbid conditions. Evidence consistently identifies delayed presentation as a major determinant of bowel resection and adverse postoperative outcomes.

The integration of Taxis-based management algorithms and laparoscopic surgical techniques offers promising opportunities for improving patient care. However, successful implementation requires careful patient selection and adherence to evidence-based protocols. The findings support the development of comprehensive risk stratification frameworks that combine epidemiological determinants with clinical decision-making processes.

Future research should focus on prospective multicenter studies, standardized reporting systems, and predictive modeling approaches capable of improving early identification of high-risk patients. Such efforts may contribute to reducing complications, optimizing surgical outcomes, and enhancing overall quality of care in adult inguinal hernia incarceration management.

## References

1. Akay T, Akici M. Should a laparoscopic repair be the first choice in incarcerated inguinal hernia. *Int Surg J.* 2020 Mar 26;7(4):961-5. doi: 10.18203/2349-2902.isj20201014.
2. Chen P, Huang L, Yang W, et al. Risk factors for bowel resection among patients with incarcerated groin hernias: a meta-analysis. *Am J Emerg Med.* 2020 Feb 1;38(2):376-83. doi: 10.1016/j.ajem.2019.09.023
3. East B, Pawlak M, de Beaux AC, et al. A manual reduction of hernia under analgesia/sedation (Taxis) in the acute inguinal hernia: a useful technique in COVID-19 times to reduce the need for emergency surgery—a literature review. *Hernia.* 2020 Oct;24:937-41. doi: 10.1007/s10029-020-02227-1.
4. Ferranti F, et al. Incarcerated Inguinal Hernia in the Elderly: InTech Open, Surgical Implications. doi: 10.5772/intechopen.1001884.
5. Merani S, Aufhauser DD, Maskin AT, et al. Not as rare as initially described: Transplant ureter incarceration within inguinal hernia. Two cases, literature review, and management algorithm. *Transplant Proc.* 2021 Sep;53(7):2285-2290. doi: 10.1016/j.transproceed.2021.07.012.
6. Pawlak M, East B, De Beaux AC, et al. Algorithm for management of an incarcerated inguinal hernia in the emergency settings with manual reduction. Taxis, the technique and its safety. *Hernia.* 2021 Oct;25(5):1253-8. doi: 10.1007/s10029-021-02429-1.
7. Sabra H, Alimoradi M, El-Helou E, et al. Perforated sigmoid colon cancer presenting as an incarcerated inguinal hernia: A case report. *Int J Surg Case Rep.* 2020 Jan 1;72:108-11. doi: 10.1016/j.ijscr.2020.05.067.
8. Sartori A, Balla A, Botteri E, et al. Laparoscopic approach in emergency for the treatment of acute incarcerated groin hernia: a systematic review and meta-analysis. *Hernia.* 2023 Jun;27(3):485-501. doi: 10.1007/s10029-022-02631-9.