



OPEN ACCESS

SUBMITTED 03 July 2025

ACCEPTED 02 August 2025

PUBLISHED 01 September 2025

VOLUME Vol.07 Issue09 2025

CITATION

COPYRIGHT

© 2025 Original content from this work may be used under the terms of the creative commons attributes 4.0 License.

Pathological And Molecular Characterization Of A Dual Bacterial Co-Infection By *Klebsiella Pneumoniae* And *Pseudomonas* *Aeruginosa* In Juvenile European Goldfinches (*Carduelis Carduelis*)

Dr. Alessandra Vitale

Department of Veterinary Medicine, University of Bologna, Italy

Abstract: The present study reports the pathological and molecular characterization of a rare dual bacterial co-infection caused by *Klebsiella pneumoniae* and *Pseudomonas aeruginosa* in juvenile European goldfinches (*Carduelis carduelis*). Affected birds presented with clinical signs of respiratory distress, lethargy, anorexia, and progressive weight loss. Necropsy revealed severe pneumonia, hepatomegaly, and splenomegaly, while histopathological analysis confirmed extensive inflammatory infiltration and tissue necrosis in the lungs and liver. Bacteriological culture yielded both *K. pneumoniae* and *P. aeruginosa*, which were further confirmed by molecular identification using species-specific PCR assays. Antimicrobial susceptibility testing demonstrated multidrug resistance, highlighting the therapeutic challenges posed by these opportunistic pathogens. The dual infection likely exacerbated disease severity, contributing to rapid mortality in the flock. This case underscores the significance of considering mixed bacterial infections in avian pathology and emphasizes the need for accurate molecular diagnostics and prudent antimicrobial stewardship in avian medicine. Furthermore, the findings provide insights into the pathogenic synergy between *K. pneumoniae* and *P. aeruginosa* in passerine birds, with implications for wildlife health surveillance and conservation.

Keywords: *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, dual infection, European goldfinch, avian pathology, molecular characterization, multidrug resistance, respiratory disease, wildlife health.

Introduction: Passerine birds, which include the European goldfinch (*Carduelis carduelis*), represent a diverse order with significant ecological roles and a growing presence in the pet bird trade [1, 2, 4]. While captive passerines are often sought for their aesthetic appeal and vocalizations, they are susceptible to a wide range of infectious diseases, with bacterial pathogens being a major cause of morbidity and mortality [4, 15]. The diagnosis and management of these infections are critical for avian health and can have significant public health implications due to the zoonotic potential of many avian pathogens [3, 11].

Among the most challenging bacterial agents are opportunistic pathogens like *Klebsiella pneumoniae* and *Pseudomonas aeruginosa*. Both are Gram-negative bacteria with a global distribution that are commonly associated with environmental contamination and can colonize various animal species, including poultry and pet birds [5, 24]. *Klebsiella pneumoniae* is a known cause of respiratory disease, yolk sac infections, and septicemia in several avian species [25, 26, 27], while *Pseudomonas aeruginosa* is notorious for causing necrotizing lesions, septicemia, and respiratory granulomas [28, 29]. The increasing prevalence of antimicrobial resistance in these pathogens is a serious concern, as it limits treatment options and facilitates the transfer of resistance genes between different bacterial populations, creating a significant "One Health" challenge [5, 6, 7].

The complexity of bacterial infections is further compounded by the phenomenon of co-infection, where two or more pathogenic agents simultaneously infect a host [33, 40]. The synergistic interactions between these pathogens can lead to more severe disease outcomes, enhanced virulence, and increased resistance to host defenses and antibiotic treatments [41, 42]. In co-infections, the presence of one pathogen can disrupt the host's normal gut microbiota, facilitating colonization by a second pathogen [16, 17, 18, 19, 20]. Additionally, interspecies communication mechanisms, such as quorum sensing, can promote biofilm formation and enhance antibiotic tolerance, complicating therapeutic interventions [43, 44, 45]. To date, reports of dual infections by *K. pneumoniae* and *P. aeruginosa* in passerine birds are rare.

This study aims to describe the pathological findings and molecular characteristics of a fatal dual co-

infection with *K. pneumoniae* and *P. aeruginosa* in a cohort of nestling European goldfinches. By providing a detailed account of this case, we hope to improve diagnostic awareness and contribute to a better understanding of the pathology and clinical significance of such co-infections in avian medicine.

Methods

Case Description and Sample Collection

Three deceased nestling European goldfinches (approximately 10 days old) were submitted to a veterinary pathology laboratory with a history of progressive lethargy, respiratory distress, and emaciation. The nestlings were part of a clutch found in a domestic setting. A full necropsy was performed immediately upon arrival to assess gross pathological changes [8, 21].

Necropsy and Histopathology

During post-mortem examination, a detailed record of all gross pathological findings was made. Tissue samples from major organs, including the liver, spleen, lungs, and kidneys, were collected, immediately fixed in 10% neutral-buffered formalin, and embedded in paraffin [8]. Sections were cut at 4 μ m and stained with hematoxylin and eosin (H&E) for histopathological examination under a light microscope. Special stains were not used.

Bacteriological Isolation and Identification

Swabs were collected aseptically from the liver, lungs, and yolk sac of each nestling. The swabs were inoculated onto various bacteriological media, including blood agar, MacConkey agar, and specific selective media for *Pseudomonas* species. The plates were incubated aerobically at 37 °C for 24–48 hours. Morphologically distinct colonies were subcultured for pure isolation and subjected to standard biochemical tests for preliminary identification [15, 21].

Molecular Characterization

To confirm the bacterial species and rule out co-infection with common avian viruses, molecular analysis was performed. Bacterial DNA was extracted from the isolated pure cultures using a commercial DNA purification kit. Polymerase Chain Reaction (PCR) assays were conducted to amplify species-specific gene targets for both *K. pneumoniae* and *P. aeruginosa* [30, 31, 32]. For *K. pneumoniae*, the *rpoB* gene was targeted, and for *P. aeruginosa*, the *ecfX* gene was utilized [31, 32]. In addition, reverse transcription PCR (RT-PCR) and real-time PCR assays were performed on tissue homogenates to test for common avian viral pathogens, including Avian avulavirus 1, Avian Bornavirus, Avian Polyomavirus, and Circovirus, to exclude a viral etiology [34, 35, 36, 37].

Antimicrobial Susceptibility Testing

Antimicrobial susceptibility testing was performed on the isolated bacterial strains using the standard Kirby-Bauer disk diffusion method according to the Clinical and Laboratory Standards Institute (CLSI) guidelines. A panel of antibiotics commonly used in veterinary medicine was tested to determine the resistance profile of each isolate [5, 22].

Results

Gross Pathological Findings

Necropsy revealed severe, multifocal to coalescing fibrinous inflammation of the air sacs, consistent with airsacculitis. The lungs were congested and edematous, with areas of consolidation. The liver appeared swollen, with multiple discrete white-to-yellow granulomatous foci visible on the surface and within the parenchyma. The spleen was moderately enlarged (splenomegaly), and the kidneys were pale with petechial hemorrhages. One of the nestlings also presented with an infected, unabsorbed yolk sac containing a caseous exudate.

Histopathological Findings

Microscopic examination of the lungs showed severe heterophilic pneumonia and marked airsacculitis, with infiltration of heterophils and macrophages. The alveolar septa were thickened, and numerous bacterial colonies were observed within the air spaces. The most striking finding in the liver was the presence of multiple, well-demarcated granulomas, characterized by a central core of necrotic debris and bacterial colonies, surrounded by a cuff of epithelioid macrophages and multinucleated giant cells. In some areas, these granulomas coalesced to form larger lesions. The spleen showed marked lymphoid depletion and diffuse histiocytic proliferation.

Bacteriological and Molecular Findings

Bacteriological cultures from the lungs, liver, and yolk sac consistently yielded both *Klebsiella pneumoniae* and *Pseudomonas aeruginosa*. The two species were identified based on colony morphology, biochemical tests, and confirmed by molecular methods. PCR assays targeting species-specific genes (*rpoB* and *ecfX*) confirmed the presence of both organisms in all tested samples [31, 32]. All tissue samples tested negative for Avian avulavirus, Avian Bornavirus, Polyomavirus, and Circovirus [34, 35, 36, 37].

Antimicrobial Susceptibility Profile

The isolated strains of *K. pneumoniae* and *P. aeruginosa* demonstrated a multidrug-resistant profile. Both strains were resistant to several broad-spectrum antibiotics, including ampicillin, tetracycline, and chloramphenicol, which are commonly used in

avian medicine. The isolates showed some susceptibility to aminoglycosides (e.g., amikacin) and fluoroquinolones (e.g., enrofloxacin), although the resistance patterns varied slightly between the two bacterial species.

Discussion

This report documents a fatal case of co-infection by *K. pneumoniae* and *P. aeruginosa* in nestling European goldfinches, providing the first detailed pathological findings of this dual infection in this species. The severity of the lesions, including granulomatous hepatitis and fibrinous airsacculitis, highlights the significant synergistic pathogenicity of these two organisms [33, 40]. The co-infection likely exacerbated the pathological outcome, as has been observed in other animal models and in-vitro studies [41, 42]. The presence of both organisms in multiple organs, including the yolk sac, suggests a systemic, septicemic process [27, 28].

The probable route of transmission in this case was environmental or parental [46, 47]. Nestlings are particularly vulnerable to infections from their immediate environment, which can be a reservoir for opportunistic bacteria [48]. Contamination of eggs or nests with fecal material from parents or other birds could have led to the initial infection [12, 13, 14, 47]. The avian gut microbiota plays a crucial role in preventing colonization by pathogens, but stress, poor sanitation, and the underdeveloped immune system of nestlings can disrupt this balance, allowing opportunistic pathogens to thrive [17, 18, 19]. The presence of the unabsorbed yolk sac provided an ideal growth medium for the bacteria, likely serving as a source of systemic infection.

The histopathological findings were consistent with the known pathology of these agents. The granulomas in the liver are a hallmark of chronic inflammatory reactions to pathogens like *Klebsiella* and are a common finding in avian bacterial diseases [15, 27]. Conversely, the extensive necrotizing and purulent inflammation in the lungs and air sacs is characteristic of *Pseudomonas* infection [38, 39, 53]. This dual presentation suggests a complex pathological process where both bacteria contributed to the overall disease picture. The molecular identification of both species confirms their roles as co-infecting agents, and the negative results for viral pathogens emphasize the primary bacterial nature of this outbreak.

The finding of multidrug resistance in both bacterial isolates is particularly concerning, underscoring the broader issue of antimicrobial resistance in wildlife and captive birds [5, 6, 7, 24]. Pet and wild birds can act as sentinels and reservoirs for resistant bacteria, which can be transmitted to humans [3, 11]. The use of molecular

techniques for rapid and accurate diagnosis is essential for guiding effective treatment and implementing biosecurity measures to prevent the spread of such resistant strains [49, 50, 51, 52]. This case highlights the need for continued surveillance of bacterial pathogens in passerine birds and a proactive "One Health" approach to managing antimicrobial resistance [11].

Conclusion

In conclusion, this study documents the pathological and molecular findings of a fatal dual infection by *Klebsiella pneumoniae* and *Pseudomonas aeruginosa* in European goldfinch nestlings. The severity of the pathological lesions and the multidrug-resistant nature of the isolates underscore the significant threat posed by these opportunistic pathogens, especially in co-infection scenarios. This report serves as a reminder of the importance of detailed diagnostic investigations in avian pathology and the need for a comprehensive understanding of co-infection dynamics to improve clinical outcomes in avian patients and protect public health.

References

1. Dorrestein, G.M. Passerines. In Handbook of Avian Medicine, 2nd ed.; Tully, T.N., Jr., Dorrestein, G.M., Jones, A.K., Eds.; Saunders: Seneca Falls, NY, USA, 2009; pp. 169–208. ISBN 9780702028748.
2. Zuccon, D.; Prys-Jones, R.; Rasmussen, P.C.; Ericson, P.G.P. The phylogenetic relationships and generic limits of finches (Fringillidae). *Mol. Phylogenet. Evol.* 2012, 62, 581–596.
3. Boseret, G.; Losson, B.; Mainil, J.G.; Thiry, E.; Saegerman, C. Zoonoses in pet birds: Review and perspectives. *Vet. Res.* 2013, 44, 36.
4. Joseph, V. Infectious and parasitic diseases of captive passerines. *Semin. Avian Exot. Pet. Med.* 2003, 12, 21–28.
5. Marouf, S.; Li, X.; Salem, H.M.; Ahmed, Z.S.; Nader, S.M.; Shaalan, M.; Awad, F.H.; Zhou, H.; Cheang, T. Molecular detection of multidrug-resistant *Pseudomonas aeruginosa* of different avian sources with pathogenicity testing and in vitro evaluation of antibacterial efficacy of silver nanoparticles against multidrug-resistant *P. aeruginosa*. *Poult. Sci.* 2023, 102, 102995.
6. Shoaib, M.; Tang, M.; Aqib, A.I.; Zhang, X.; Wu, Z.; Wen, Y.; Hou, X.; Xu, J.; Hao, R.; Wang, S.; et al. Dairy farm waste: A potential reservoir of diverse antibiotic resistance and virulence genes in aminoglycoside—And beta-lactam-resistant *Escherichia coli* in Gansu Province, China. *Environ. Res.* 2024, 263, 120190.
7. Asokan, S.; Jacob, T.; Jacob, J.; AlSosowaa, A.A.; Cherian, T.; Peijnenburg, W.J.G.M.; Vijayan, S. *Klebsiella pneumoniae*: A growing threat in the era of antimicrobial resistance. *Microbe* 2025, 7, 100333.
8. Schmidt, R.E.; Reavill, D.R.; Phalen, D.N. *Pathology of Pet and Aviary Birds*, 2nd ed.; Wiley Blackwell: Hoboken, NJ, USA, 2015; pp. 95–125.
9. Nolan, L.K.; Vaillancourt, J.P.; Barbieri, N.L.; Louge, C.M. Colibacillosis. In *Diseases of Poultry*; Swayne, D.E., Ed.; Wiley Online Library: Ames, IA, USA, 2020; pp. 770–830.
10. Qui, Y.; Lv, C.; Chen, J.; Sun, Y.; Tang, T.; Zhang, Y.; Yang, Y.; Wang, G.; Xu, Q.; Zhang, X.; et al. The global distribution and diversity of wild-bird-associated pathogens: An integrated data analysis and modeling study. *Med* 2025, 6, 100553.
11. Kobuszewska, A.; Wysok, B. Pathogenic Bacteria in Free-Living Birds, and Its Public Health Significance. *Animals* 2024, 14, 968.
12. Gantois, I.; Ducatelle, R.; Pasmans, F.; Haesebrouck, F.; Gast, R.; Humphrey, T.J.; Van Immerseel, F. Mechanisms of egg contamination by *Salmonella enteritidis*. *FEMS Microbiol. Rev.* 2009, 33, 718–738.
13. Gast, R.K.; Porter, R.E., Jr. *Salmonella Infections*. In *Diseases of Poultry*, 14th ed.; Swayne, D.E., Boulianne, M., Logue, S.M., Eds.; Wiley: Hoboken, NJ, USA, 2020; pp. 719–753.
14. Ottinger, M.A.; Mani, S.; Krüger, S.; Coverdale, B.; Willows-Munro, S.; Combrink, L. Microbiomes in Birds: A review of links to health and reproduction. *Reprod. Fertil.* 2024, 5, e230076.
15. Gerlach, H. Bacteria. In *Avian Medicine: Principles and Application*; Ritchie, B.W., Harrison, G.J., Harrison, L.R., Eds.; Wingers Publishing, Inc.: Lake Worth, FL, USA, 1994; pp. 949–983.
16. Zhou, Y.; Li, Y.; Zhang, L.; Wu, Z.; Huang, Y.; Yan, H.; Zhong, J.; Wang, L.J.; Abdullah, H.M.; Wang, H.H. Antibiotic administration routes and oral exposure to antibiotic resistant bacteria as key drivers for gut microbiota disruption and resistome in poultry. *Front. Microbiol.* 2020, 11, 1319.
17. Sun, F.; Chen, J.; Liu, K.; Tang, M.; Yang, Y. The avian gut microbiota: Diversity, influencing factors, and future directions. *Front. Microbiol.* 2022, 13, 934272.
18. Benskin, C.M.H.; Rhodes, G.; Pickup, R.W.; Wilson, K.; Hartley, I.R. Diversity and temporal stability of bacterial communities in a model passerine bird, the zebra finch. *Mol. Ecol.* 2010, 19, 5531–5544.
19. Garcia-Mazcorro, J.F.; Castillo-Carranza, S.A.; Guard, B.; Gomez-Vazquez, J.P.; Dowd, S.E.; Brightsmith, D.J. Comprehensive molecular characterization of

bacterial communities in feces of bet birds using 16S marker sequencing. *Microb. Ecol.* 2017, 73, 224–235.

20. Matteucci, G.; Binanti, D. Presence and Significance of Gram-negative bacteria in the cloacal microbiota of Psittacine and Passerine. In *Proceedings of the ICARE 2017—International Conference of Avian Herpetological and Exotic Mammal Medicine*, Venice, Italy, 25–29 March 2017.

21. Coles, B.H.; Krautwald-Junghanns, M.; Orosz, S.E.; Tully, T. *Essentials of Avian Medicine and Surgery*, 3rd ed.; Wiley-Blackwell: Hoboken, NJ, USA, 2007.

22. El Fertas-Aissani, R.; Messai, Y.; Alouache, S.; Bakour, R. Virulence profiles and antibiotic susceptibility patterns of *Klebsiella pneumoniae* strains isolated from different clinical specimens. *Pathol. Biol.* 2013, 61, 209–216.

23. Davies, Y.M.; Cunha, M.P.V.; Oliveira, M.G.X.; Oliveira, M.C.V.; Philadelpho, N.; Romero, D.C.; Milanelo, L.; Guimarães, M.B.; Ferreira, A.J.P.; Moreno, A.M.; et al. Virulence and antimicrobial resistance of *Klebsiella pneumoniae* isolated from passerine and psittacine birds. *Avian Pathol.* 2016, 45, 194–201.

24. Ahmed, H.A.; Awad, N.F.S.; Abd El-Hamid, M.I.; Shaker, A.; Mohamed, R.E.; Elsohaby, I. Pet birds as potential reservoir of virulent and antibiotic resistant zoonotic bacteria. *Comp. Immunol. Microbiol. Infect. Dis.* 2021, 75, 101606.

25. Razmyar, J.; Zamani, A.H. An outbreak of yolk sac infection and dead-in-shell mortality in common canary (*Serinus canaria*) caused by *Klebsiella pneumoniae*. *Iran. J. Vet. Res.* 2016, 17, 141–143.

26. Nakhaee, P.; Zarif Moghadam, H.; Shokrpour, S.; Razmyar, J. *Klebsiella pneumoniae* infection in canaries (*Serinus canaria domestica*): A case report. *Iran. J. Vet. Res.* 2022, 23, 280–284.

27. Kock, N.; Kock, R. Disease epidemic in lesser flamingos (*Phoeniconauis minor*) in Kenya. In *Proceedings of the American Association of Zoo Veterinarians; American Association of Zoo Veterinarians*: East Lansing, MI, USA, 1995; p. 115.

28. Momotani, E.; Kiryu, M.; Ohshiro, M.; Murakami, Y.; Ashida, Y.; Watanabe, S.; Matsubara, Y. Granulomatous lesions caused by *Pseudomonas aeruginosa* in the Ostrich (*Struthio camelus*). *J. Comp. Pathol.* 1995, 112, 273–282.

29. Bailey, T.A.; Silvanose, C.D.; Naldo, J.N.; Howlett, J.H. *Pseudomonas aeruginosa* infections in kori bustards (*Ardeotis kori*). *Avian Pathol.* 2000, 29, 41–44.

30. Diancourt, L.; Passet, V.; Verhoef, J.; Grimont, P.A.D.; Brisse, S. Multilocus sequence typing of

Klebsiella pneumoniae nosocomial isolates. *J. Clin. Microbiol.* 2005, 43, 4178–4182.

31. Lavenir, R.; Jocktane, D.; Laurent, F.; Nazaret, S.; Cournoyer, B. Improved reliability of *Pseudomonas aeruginosa* PCR detection by the use of the species-specific *ecfX* gene target. *J. Microbiol. Methods* 2007, 70, 20–29.

32. He, Y.; Guo, X.; Xiang, S.; Li, J.; Li, X.; Xiang, H.; He, J.; Chen, D.; Chen, J. Comparative analyses of phenotypic methods and 16S rRNA, *khe*, *rpoB* genes sequencing for the identification of clinical isolates of *Klebsiella pneumoniae*. *Antonie Van Leeuwenhoek* 2016, 109, 1029–1040.

33. Palazzolo, S.; Gervasi, C.; Abbate, J.M.; Gjurčević, E.; Falletti, R.; Piro, M.G.; Lanteri, G.; Iaria, C.; Marino, F. Natural bacterial co-infection in farmed European sea bass intended for experimental research in Sicily, southern Italy: Pathological findings. *Fishes* 2024, 9, 360.

34. Sutton, D.A.; Allen, D.P.; Fuller, C.M.; Mayers, J.; Mollett, B.C.; Londt, B.Z.; Reid, S.M.; Mansfield, K.L.; Brown, I.H. Development of an avian avulavirus 1 (AAvV-1) L-gene real-time RT-PCR assay using minor groove binding probes for application as a routine diagnostic tool. *J. Virol. Methods* 2019, 265, 9–14.

35. Sigrist, B.; Geers, J.; Albini, S.; Rubbenstroth, D.; Wolfrum, N. A New Multiplex Real-Time RT-PCR for Simultaneous Detection and Differentiation of Avian Bornaviruses. *Viruses* 2021, 13, 1358.

36. Johne, R.; Enderlein, D.; Nieper, H.; Müller, H. Novel polyomavirus detected in the feces of a chimpanzee by nested broad-spectrum PCR. *J. Virol.* 2005, 79, 3883–3887.

37. Todd, D.; Weston, J.; Ball, N.W.; Borghmans, B.J.; Smyth, J.A.; Gelmini, L.; Lavazza, A. Nucleotide sequence-based identification of a novel circovirus of canaries. *Avian Pathol.* 2001, 30, 321–325.

38. Okada, F.; Ando, Y.; Honda, K.; Nakayama, T.; Ono, A.; Tanoue, S.; Maeda, T.; Mori, H. Acute *Klebsiella pneumoniae* pneumonia alone and with concurrent infection: Comparison of clinical and thin-section CT findings. *Br. J. Radiol.* 2010, 83, 854–860.

39. Abd El-Ghany, W.A. *Pseudomonas aeruginosa* infection of avian origin: Zoonosis and one health implications. *Vet. World* 2021, 14, 2155–2159.

40. Jones-Nelson, O.; Hilliard, J.J.; DiGiandomenico, A.; Warrenner, P.; Alfaro, A.; Cheng, L.; Stover, C.K.; Cohen, T.S.; Sellman, B.R. The Neutrophilic response to *Pseudomonas* damages the airway barrier, promoting infection by *Klebsiella pneumoniae*. *Am. J. Respir. Cell Mol. Biol.* 2018, 59, 745–756.

41. Marchaim, D.; Perez, F.; Lee, J.; Bheemreddy, S.; Hujer, A.M.; Rudin, S.; Hayakawa, K.; Lephart, P.R.; Blunden, C.; Shango, M.; et al. "Swimming in resistance": Co-colonization with carbapenem-resistant Enterobacteriaceae and *Acinetobacter baumannii* or *Pseudomonas aeruginosa*. *Am. J. Infect. Control* 2012, 40, 830–835.
42. Muzaki, M.Z.B.M.; Subramoni, S.; Summers, S.; Kjelleberg, S.; Rice, S.A. *Klebsiella pneumoniae* AI-2 transporters mediate interspecies interactions and composition in a three-species biofilm community. *NPJ Biofilms Microbiomes* 2024, 10, 91.
43. Elias, S.; Banin, E. Multi-species biofilms: Living with friendly neighbors. *FEMS Microbiol. Rev.* 2012, 36, 990–1004.
44. Castillo-Juárez, I.; Maeda, T.; Mandujano-Tinoco, E.A.; Tomás, M.; Pérez-Eretza, B.; García-Contreras, S.J.; Wood, T.K.; García-Contreras, R. Role of quorum sensing in bacterial infections. *World J. Clin. Cases* 2015, 3, 575–598.
45. Cui, S.; Kim, E. Quorum sensing and antibiotic resistance in polymicrobial infections. *Commun. Integr. Biol.* 2024, 17, 2415598.
46. Martin-Vivaldi, M.; Soler, J.J.; Peralta-Sánchez, J.M.; Arco, L.; Martín-Platero, A.M.; Martínez-Bueno, M.; Ruiz-Rodríguez, M.; Valdivia, E. Special structures of hoopoe eggshells enhance the adhesion of symbiont-carrying uropygial secretion that increase hatching success. *J. Anim. Ecol.* 2014, 83, 1289–1301.
47. Van Veelen, H.P.J.; Salles, J.F.; Tieleman, B.I. Microbiome assembly of avian eggshells and their potential as transgenerational carriers of maternal microbiota. *ISME J.* 2018, 12, 1375.
48. Chen, C.Y.; Chen, C.K.; Chen, Y.Y.; Fang, A.; Shaw, G.T.W.; Hung, C.M.; Wang, D. Maternal gut microbes shape the early-life assembly of gut microbiota in passerine chicks via nests. *Microbiome* 2020, 8, 129.
49. Gibbs, P.S.; Kasa, R.; Newbrey, J.L.; Petermann, S.R.; Wooley, R.E.; Winson, H.M.; Reed, W. Identification, antimicrobial resistance profiles and virulence of members from the family Enterobacteriaceae from the feces of yellow-headed blackbirds (*Xanthocephalus xanthocephalus*) in North Dakota. *Avian Dis.* 2007, 51, 649–655.
50. Horn, R.V.; Cardoso, E.M.; Lopes, E.S.; Teixeira, R.S.C.; Albuquerque, A.H.; Rocha-e-Silva, R.C.; Machado, D.N.; Bezerra, W.G.A. Identification and antimicrobial resistance of members from the Enterobacteriaceae family isolated from canaries (*Serinus canaria*). *Pesq. Vet. Bras.* 2015, 35, 552–556.
51. Stenkat, J.; Krautwald-Junghanns, M.E.; Schmitz, O.A.; Eilers, A.; Schmidt, V. Aerobic cloacal and pharyngeal bacterial flora in six species of free-living birds. *J. Appl. Microbiol.* 2014, 117, 1564–1571.
52. Raue, R.; Schmidt, V.; Freick, M.; Reinhardt, B.; Johne, R.; Kamphausen, L.; Kaleta, E.F.; Muller, H.; Krautwald-Junghanns, M.E. A disease complex associated with pigeon circovirus infection, young pigeon disease syndrome. *Avian Pathol.* 2005, 34, 418–425.
53. Jackson, N.K.; Philips, S.N. Necrotizing hepatitis in pet birds associated with *Pseudomonas fluorescens*. *Avian Dis.* 1996, 40, 473–476.